

Differential Geometry of the Lie algebra of the quantum plane

Salih Çelik ¹ and Sultan A Çelik

Yildiz Technical University, Department of Mathematics,
34210 Davutpasa-Esenler, Istanbul, TURKEY.

We present a differential calculus on the extension of the quantum plane obtained considering that the (bosonic) generator x is invertible and furthermore working polynomials in $\ln x$ instead of polynomials in x . We call quantum Lie algebra to this extension and we obtain its Hopf algebra structure and its dual Hopf algebra.

¹E-mail address: sacelik@yildiz.edu.tr

Quantum plane [1] is a simple example of quantum space and has been studied intensively by many authors in the past years. It can be obtained by deformation of the classical plane [2]. For references to the literature we refer to the recent book by Majid [3].

As usual in noncommutative geometry [4] quantum plane has many differential calculi. By using the noncommutative differential geometry method in Ref. 5 and interpreting the dual plane of the quantum plane to consist of differentials of the coordinates of the quantum plane, covariant differential calculus on the quantum plane has been developed. Therefore the quantum plane provides a simple example for noncommutative differential geometry [6].

It is known that, in order to construct a noncommutative differential calculus on quantum groups and Hopf algebras, one takes into consideration the associative algebra of functions on the group. The starting point of this work is its Lie algebra. For quantum superplane, similar a work studied in Ref. 7. In this work, we present here a differential calculus on the Lie algebra of the associative algebra of functions on the extended quantum plane.

The quantum plane [1] is defined as an associative algebra generated by two noncommuting coordinates x and y with the single quadratic relation

$$xy - qyx = 0 \quad q \in \mathbb{C} - \{0\}. \quad (1)$$

This algebra is known as the algebra of polynomials over the quantum plane and we shall denote by $\mathcal{A}_q = Fun_q(R(2))$. We extend the algebra $\mathcal{A}_q$ by including inverse of x which obeys

$$xx^{-1} = 1 = x^{-1}x.$$

We know, from Ref. 8, that the extended algebra $\mathcal{A}_q$ is a Hopf algebra with the following costructures:

$$\begin{aligned} \Delta(x) &= x \otimes x & \Delta(y) &= y \otimes 1 + x \otimes y \\ \epsilon(x) &= 1 & \epsilon(y) &= 0 \\ \kappa(x) &= x^{-1} & \kappa(y) &= -x^{-1}y. \end{aligned} \quad (2)$$

It is not difficult to verify the following properties of costructures:

$$(\Delta \otimes \text{id}) \circ \Delta = (\text{id} \otimes \Delta) \circ \Delta$$

$$m \circ (\epsilon \otimes \text{id}) \circ \Delta = m \circ (\text{id} \otimes \epsilon) \circ \Delta \quad (3)$$

$$m \circ (\kappa \otimes \text{id}) \circ \Delta = \epsilon = m \circ (\text{id} \otimes \kappa) \circ \Delta$$

where id denotes the identity mapping and m is the multiplication map

$$m : \mathcal{A}_q \otimes \mathcal{A}_q \longrightarrow \mathcal{A}_q \quad m(a \otimes b) = ab.$$

It is known that an element of a Lie group can be represented by exponential of an element of its Lie algebra. Using this fact, we can define the generators of $\mathcal{A}_q$ as [9]

$$x = e^X \quad y = e^X Y \quad q = e^h. \quad (4)$$

Then we obtain the relation

$$[X, Y] = hY \quad (5)$$

where

$$[a, b]_{\pm} = ab \pm ba.$$

This is the relations of a Lie algebra and we shall denote it by $\mathcal{L}(\mathcal{A}_q) =: g_h$. The Hopf algebra structure of g_h can be read off from (2),

$$\begin{aligned} \Delta(X) &= X \otimes 1 + 1 \otimes X \\ \Delta(Y) &= Y \otimes e^{-X} + 1 \otimes Y \\ \epsilon(X) &= 0 \quad \epsilon(Y) = 0 \\ \kappa(X) &= -X \quad \kappa(Y) = -Y e^X. \end{aligned} \quad (6)$$

We now shall build up a noncommutative differential calculus on the Lie algebra g_h . This must be involve functions on the Lie algebra g_h , differentials and differential forms. For this, we have to define a linear map δ which acts on the functions of the elements of g_h . The exterior differential δ is an operator which gives the mapping from the generators of g_h to the differentials:

$$\delta : u \mapsto \delta u \quad u \in \{X, Y\}.$$

We demand that the exterior differential δ has to satisfy two properties: the nilpotency

$$\delta^2 = 0 \quad (7a)$$

and the Leibniz rule

$$\delta(fg) = (\delta f)g + f(\delta g). \quad (7b)$$

In order to establish a noncommutative differential calculus on the algebra g_h , we assume that the commutation relations between the elements of g_h and their differentials are the following form:

$$\begin{aligned} X \delta X &= A_{11} \delta X X + B_1 \delta X + B_2 \delta Y \\ X \delta Y &= A_{12} \delta Y X + B_3 \delta X + B_4 \delta Y \\ Y \delta X &= A_{21} \delta X Y + B_5 \delta X + B_6 \delta Y \\ Y \delta Y &= A_{22} \delta Y Y + B_7 \delta X + B_8 \delta Y. \end{aligned} \quad (8)$$

The coefficients A_{ij} and B_i will be determined in terms of the "new" deformation parameter h . To find them we shall use the consistency of calculus.

Using the consistency conditions

$$\begin{aligned} \delta(XY - YX - hY) &= 0 \\ (XY - YX - hY)\delta X &= 0 \\ (XY - YX - hY)\delta Y &= 0 \end{aligned}$$

we obtain the system

$$\begin{aligned} A_{11} &= A_{12} = A_{21} = 0 & B_2(1 - A_{22}) &= 0 \\ B_3 - B_5 &= 0 & B_4 - B_6 &= h & B_3B_6 - B_2B_7 &= hB_5 \\ (B_1 - B_4)B_6 &+ (B_8 - B_5)B_2 &= -hB_6 \\ B_3B_6 - B_2B_7 &= hB_8 \\ (B_1 - B_4)B_7 &+ (B_8 - B_5)B_3 &= hB_7 \end{aligned}$$

or

$$\begin{aligned} A_{11} &= A_{12} = A_{21} = 0 & B_2(1 - A_{22}) &= 0 \\ B_3 &= B_5 = B_8 & B_4 - B_6 &= h & (B_1 - B_4 - h)B_7 &= 0. \end{aligned}$$

Thus we have the following algebra

$$[X, \delta X] = B_1 \delta X + B_2 \delta Y$$

$$\begin{aligned}
[X, \delta Y] &= B_3 \delta X + B_4 \delta Y \\
[Y, \delta X] &= B_3 \delta X + (B_4 - h) \delta Y \\
[Y, \delta Y] &= (A_{22} - 1) \delta Y Y + B_7 \delta X + B_3 \delta Y.
\end{aligned} \tag{9}$$

But we do not have the required relations because of the constants. To find the constants in the above relations, we shall consider the covariance of the noncommutative differential calculus.

We first note that consistency of a differential calculus with commutation relation (5) means that the algebra is a graded associative algebra generated by the elements of the set $\{X, Y, \delta X, \delta Y\}$. So, it is sufficient only describe the actions of co-maps on the subset $\{\delta X, \delta Y\}$.

Let Γ be a free left module over the algebra g_h generated by the elements of the set $\{X, Y, \delta X, \delta Y\}$. The module Γ becomes a unital associative algebra if one defines a multiplication law on Γ by relations (5) and (9).

We consider a map $\phi_L : \Gamma \longrightarrow g_h \otimes \Gamma$ such that

$$\phi_L \circ \delta = (\text{id} \otimes \delta) \circ \Delta. \tag{10}$$

Thus we have

$$\phi_L(\delta X) = 1 \otimes \delta X \quad \phi_L(\delta Y) = 1 \otimes \delta Y + \frac{1 - e^h}{h} Y \otimes X e^{-X}. \tag{11}$$

We now define a map Δ_L as follows:

$$\Delta_L(a_1 \delta b_1 + \delta b_2 a_2) = \Delta(a_1) \phi_L(\delta b_1) + \phi_L(\delta b_2) \Delta(a_2). \tag{12}$$

Applying the linear map Δ_L to relations (9) we get, as the solution of the obtained system,

$$\begin{aligned}
A_{11} = A_{12} = A_{21} = 1 \quad A_{22} = e^h \\
B_1 = B_6 = -h \quad B_2 = B_3 = B_4 = B_5 = B_7 = B_8 = 0.
\end{aligned} \tag{13}$$

Consequently, the relations (9) is of the form

$$\begin{aligned}
[X, \delta X] &= -h \delta X \quad [X, \delta Y] = 0 \\
[Y, \delta X] &= -h \delta Y \quad [Y, \delta Y] = (e^h - 1) \delta Y Y.
\end{aligned} \tag{14}$$

Applying the exterior differential δ to the relation (14) one has

$$[\delta X, \delta Y]_+ = 0 \quad (\delta X)^2 = 0 = (\delta Y)^2. \quad (15)$$

Note that, the differentials of X and Y in terms of x and y as follows:

$$\delta X = \frac{h}{1 - e^{-h}} \delta x x^{-1} \quad \delta Y = x^{-1}(\delta y - \delta x x^{-1}y).$$

We now consider another map $\phi_R : \Gamma \longrightarrow \Gamma \otimes g_h$ such that

$$\phi_R \circ \delta = (\delta \otimes \text{id}) \circ \Delta. \quad (16)$$

Thus one has

$$\phi_R(\delta X) = \delta X \otimes 1 \quad \phi_R(\delta Y) = \delta Y \otimes e^{-X}. \quad (17)$$

We define a map Δ_R with again (12) by replacing R with L . The map Δ_R also leaves invariant the relations (14) and (15), and the following identities are satisfied:

$$\begin{aligned} (\Delta \otimes \text{id}) \circ \Delta_L &= (\text{id} \otimes \Delta_L) \circ \Delta_L & (\epsilon \otimes \text{id}) \circ \Delta_L &= \text{id} \\ (\text{id} \otimes \Delta) \circ \Delta_R &= (\Delta_R \otimes \text{id}) \circ \Delta_R & (\text{id} \otimes \epsilon) \circ \Delta_R &= \text{id}. \end{aligned} \quad (18)$$

To denote the coproduct, counit and coinverse which will be defined on the algebra Γ with those of $\mathcal{A}$ may be inadvisable. For this reason, we shall denote them with a different notation. Let us define the map $\hat{\Delta}$ as

$$\hat{\Delta} = \Delta_R + \Delta_L \quad (19)$$

which will allow us to define the coproduct of the differential algebra. We denote the restriction of $\hat{\Delta}$ to the algebra g_h by Δ and the extension of Δ to the differential algebra Γ by $\hat{\Delta}$. It is possible to interpret the relation

$$\hat{\Delta}|_{g_h} = \Delta \quad (20)$$

as the definition of $\hat{\Delta}$ on the generators of g_h and (19) as the definition of $\hat{\Delta}$ on differentials.

It is not difficult to verify the following conditions:

a) Γ is an g_h -bimodule,

b) Γ is an g_h -bicomodule with left and right coactions Δ_R and Δ_L , respectively, making Γ a left and right g_h -comodule with (18), and

$$(\Delta_L \otimes \text{id}) \circ \Delta_R = (\text{id} \otimes \Delta_R) \circ \Delta_L \quad (21)$$

which is the g_h -bimodule property. So, the triple $(\Gamma, \Delta_L, \Delta_R)$ is a bicovariant bimodule over Hopf algebra g_h . In addition, since

c) (Γ, δ) is a first order differential calculus over g_h , and

d) δ is both a left and a right comodule map, i.e. for all $a \in g_h$

$$(\text{id} \otimes \delta) \Delta(a) = \Delta_L(\delta a) \quad (\delta \otimes \text{id}) \Delta(a) = \Delta_R(\delta a) \quad (22)$$

the quadruple $(\Gamma, d, \Delta_L, \Delta_R)$ is a first order bicovariant differential calculus over Hopf algebra g_h .

Now let us return Hopf algebra structure of Γ . If we define a counit $\hat{\epsilon}$ for the differential algebra as

$$\hat{\epsilon} \circ \delta = \delta \circ \epsilon = 0 \quad (23)$$

and

$$\hat{\epsilon}|_{g_h} = \epsilon \quad \epsilon|_{\Gamma} = \hat{\epsilon}. \quad (24)$$

we have

$$\hat{\epsilon}(\delta X) = 0 \quad \hat{\epsilon}(\delta Y) = 0 \quad (25)$$

where

$$\hat{\epsilon}(a_1 \delta b_1 + \delta b_2 a_2) = \epsilon(a_1) \hat{\epsilon}(\delta b_1) + \hat{\epsilon}(\delta b_2) \epsilon(a_2). \quad (26)$$

Here we used the fact that $\delta(1) = 0$.

As the next step we obtain a coinverse $\hat{\kappa}$. For this, it suffices to define $\hat{\kappa}$ such that

$$\hat{\kappa} \circ \delta = \delta \circ \kappa \quad (27)$$

and

$$\hat{\kappa}|_{g_h} = \kappa \quad \kappa|_{\Gamma} = \hat{\kappa} \quad (28)$$

where

$$\hat{\kappa}(a_1 \delta b_1 + \delta b_2 a_2) = \hat{\kappa}(\delta b_1) \kappa(a_1) + \kappa(a_2) \hat{\kappa}(\delta b_2). \quad (29)$$

Thus the action of $\hat{\kappa}$ on the generators δX and δY is as follows:

$$\hat{\kappa}(\delta X) = -\delta X \quad \hat{\kappa}(\delta Y) = e^{X-h} \left(\delta Y + \frac{e^h - 1}{h} \delta X Y \right). \quad (30)$$

Note that it is easy to check that $\hat{\epsilon}$ and $\hat{\kappa}$ leave invariant the relations (14) and (15). Consequently, we can say that the structure $(\Gamma, \hat{\Delta}, \hat{\epsilon}, \hat{\kappa})$ is a graded Hopf algebra.

Just as we introduced the derivatives of the generators of $\mathcal{A}_q$ in the standard way, let us introduce derivatives of the generators of g_h and multiply explicit expression of the exterior differential δ from the right by Xf and Yf , respectively. Then, using the Leibniz rule for partial derivatives

$$\partial_i(fg) = (\partial_i f)g + f(\partial_i g)$$

we obtain

$$\begin{aligned} [\partial_X, X] &= \frac{h}{1 - e^{-h}} (1 + (e^{-h} - 1)\partial_X) \\ [\partial_X, Y] &= 0 \quad [\partial_Y, X] = 0 \\ [\partial_Y, Y] &= e^h + (1 - e^h)(\partial_X - Y\partial_Y). \end{aligned} \quad (31)$$

Note that, the partial derivatives of X and Y in terms of x and y as follows:

$$\partial_X = x\partial_x + y\partial_y \quad \partial_Y = e^h x\partial_y.$$

The commutation relations among the partial derivatives can be easily obtained by using $\delta^2 = 0$. So it follows that

$$\begin{aligned} \delta &= \frac{1 - e^{-h}}{h} \delta X \partial_X + e^{-h} \delta Y \partial_Y \\ 0 &= \delta^2 = \delta \left(\frac{1 - e^{-h}}{h} \delta X \partial_X + e^{-h} \delta Y \partial_Y \right) \\ &= e^{-h} \frac{e^{-h} - 1}{h} \delta X \delta Y (\partial_Y \partial_X - \partial_X \partial_Y) \end{aligned}$$

which says that

$$[\partial_X, \partial_Y] = 0. \quad (32)$$

In order to find the commutation relations between the differentials and partial derivatives, we demand a form similar to (8). After some tedious but straightword calculations, we find

$$\begin{aligned}
[\partial_X, \delta X] &= (e^h - 1)\delta X \partial_X - e^h \delta X \\
[\partial_X, \delta Y] &= (e^h - 1)\delta Y \partial_X - e^h \delta Y \\
[\partial_Y, \delta X] &= 0 \\
[\partial_Y, \delta Y] &= (e^h - 1)\delta Y \partial_Y + \frac{(e^h - 1)^2}{h} \delta X \partial_X + e^h \frac{1 - e^h}{h} \delta X.
\end{aligned} \tag{33}$$

We now define two one-forms using the generators of g_h and show that the algebra of forms is a graded Hopf algebra.

If we call them w_1 and w_2 then we can define them as follows:

$$w_1 = \frac{1 - e^{-h}}{h} \delta X \quad w_2 = e^X \delta Y. \tag{34}$$

We denote the algebra of forms generated by w_1 and w_2 by Ω . The generators of the algebra Ω with the generators of g_h satisfy the following relations:

$$\begin{aligned}
[w_1, X] &= h w_1 & [w_1, Y] &= (1 - e^{-h}) e^{-X} w_2 \\
[w_2, X] &= 0 & [w_2, Y] &= 0.
\end{aligned} \tag{35}$$

The commutation rules of the generators of Ω are

$$w_1 w_2 = -e^h w_2 w_1 \quad w_1^2 = 0 = w_2^2. \tag{36}$$

One can make the algebra Ω into a graded Hopf algebra with the following costructures: the coproduct $\Delta : \Omega \longrightarrow \Omega \otimes \Omega$ is given by

$$\begin{aligned}
\Delta(w_1) &= w_1 \otimes 1 + 1 \otimes w_1 \\
\Delta(w_2) &= w_2 \otimes 1 + e^X \otimes w_2 - e^X Y \otimes w_1.
\end{aligned} \tag{37}$$

The counit $\epsilon : \Omega \longrightarrow \mathcal{C}$ is given by

$$\epsilon(w_1) = 0 \quad \epsilon(w_2) = 0 \tag{38}$$

and the coinverse $\kappa : \Omega \longrightarrow \Omega$ is defined by

$$\kappa(w_1) = -w_1 \quad \kappa(w_2) = e^{-X}w_2 + Yw_1. \quad (39)$$

Now we shall obtain the universal enveloping algebra of g_h , $\mathcal{U}(g_h)$, via the vector fields related to the one-forms. We first write the Cartan-Maurer forms as

$$\delta X = \frac{h}{1 - e^{-h}} w_1 \quad \delta Y = e^{-X}w_2. \quad (40)$$

Writing the exterior differential δ in the form

$$\delta = w_1 T_1 + w_2 T_2$$

and considering an arbitrary function f of the generators of g_h and using the nilpotency of δ one has

$$T_2 T_1 = e^{-h} T_1 T_2 + T_2. \quad (41)$$

The commutation relations of vector fields with the generators of g_h as follows:

$$\begin{aligned} [T_1, X] &= \frac{h}{1 - e^{-h}} (1 + (e^{-h} - 1)T_1) \\ [T_1, Y] &= 0 \quad [T_2, X] = 0 \\ [T_2, Y] &= e^{-X} (1 + (e^{-h} - 1)T_1). \end{aligned} \quad (42)$$

The Hopf algebra structure of the vector fields is given by

$$\begin{aligned} \Delta(T_1) &= T_1 \otimes 1 + 1 \otimes T_1 + (e^{-h} - 1)T_1 \otimes T_1 \\ \Delta(T_2) &= T_2 \otimes 1 + 1 \otimes T_2 + (e^{-h} - 1)T_2 \otimes T_1 \\ \epsilon(T_1) &= 0 \quad \epsilon(T_2) = 0 \\ \kappa(T_1) &= -T_1 (1 + (e^{-h} - 1)T_1)^{-1} \\ \kappa(T_2) &= -T_2 \left\{ 1 - (e^{-h} - 1)T_1 (1 + (e^{-h} - 1)T_1)^{-1} \right\}. \end{aligned} \quad (43)$$

If we introduce the operators H and N as

$$T_1 = \frac{e^{-hN} - 1}{e^{-h} - 1} \quad \text{and} \quad T_2 = e^{-hN} H \quad (44)$$

we get

$$[H, N] = H. \quad (45)$$

Then, dual Hopf algebra to $\mathcal{U}(g_h)$ is given by

$$\begin{aligned} \Delta(N) &= N \otimes 1 + 1 \otimes N \\ \Delta(H) &= H \otimes 1 + e^{hN} \otimes H \\ \epsilon(N) &= 0 \quad \epsilon(H) = 0 \\ \kappa(N) &= -N \quad \kappa(H) = -e^{-hN} H. \end{aligned} \quad (46)$$

Acknowledgment

This work was supported in part by TBTAk the Turkish Scientific and Technical Research Council.

References

- [1] Manin Yu I 1988 (Montreal Univ Preprint)
- [2] Fairlie D and Zachos C 1991 *Phys Lett B* **256** 43
- [3] Majid S 1995 "Foundations of Quantum Group Theory" (Cambridge Univ Press)
- [4] Connes A 1994 "Noncommutative Differential Geometry" (Academic Press San Diego)
- [5] Woronowicz S L 1987 *Commun Math Phys* **111** 613; 1989 *ibid* **122** 125
- [6] Wess J and Zumino B 1990 *Nucl Phys* (Proc Suppl) **18** B 302
- [7] Celik S 2002 q-alg/0201170
- [8] Brzezinski T and Majid S 1992 *Lett Math Phys* **26** 67;
Brzezinski T 1993 *Lett Math Phys* **27** 287;
Celik S A and Celik S 2000 *Int J Mod Phys A* **15** 3237
- [9] Fronsda1 C and Galindo A 1993 *Lett Math Phys* **27** 59